

**KEY INDICATOR FOR SURE START BASELINE SURVEY-2007  
KANPUR DEHAT DISTRICT OF UTTAR PARDESH**

**I. Population data of 2001 census**

Population	2,872,335
Decadal population growth rate	23.66

**II .Sample population**

Population covered	170740
Number of household covered	27805
Number of women identified with live birth in year 2007	3689
Number of women interviewed	2472

**III. Background characteristics of women interviewed**

Percentage of women below age 25	41.95
Mean age at marriage	16.95
Percentage of illiterate	55.62
Percentage of having 10+ years of schooling	31.06
Percentage of illiterate husband	24.45
Percentage of husband having 10+ years of schooling	45.86
Percentage of women able to go to health facility for taking treatment (alone)	23.30
<b>Caste</b>	
High	19.46
Middle	24.76
Low middle	29.05
Low	11.77
Muslim	12.42
Other	0.12
Hindu not specified caste	2.43
<b>Asset Index</b>	
Percentage of 1st quintile(Low)	24.11
Percentage of 2nd quintile(Low medium)	18.93
Percentage of 3rd quintile(Medium)	18.04
Percentage of 4th quintile(Medium High)	18.37
Percentage of 5th quintile(High)	20.55

**IV. Reproductive history of women**

Mean children ever born	2.12
Mean no of children surviving	1.75

**V. Mortality & Morbidity**

Neonatal Mortality rate	44.46
Infant Mortality rate	33.61
Morbidity during pregnancy (Percent )	63.71
Morbidity during labor and delivery (Percent )	33.50
Morbidity during postpartum period (Percent )	46.89
New born morbidity (Percent )	40.29

**VI. Healthy pregnancy practices**

Percentage of women received ANC during third trimester	75.0
Percentage of women received TT during pregnancy	93.33
Percentage of women received IFA tablets during pregnancy	69.54

**VII. Delivery status**

Institutional Deliveries	35.76
Home Deliveries	63.67

**VIII. Postpartum care**

Received supplementary food from AWC during postpartum period	67.31
Home visit by health worker during postpartum period	60.36

**IX. Healthy newborn practices**

Percentage of respondents practices healthy new born care	100.00
High	25.90
Intermediate	51.88
Low	22.22
Colostrum feeding	66.32
Applying nothing on the cord	49.81

**X. Awareness level**

Level of awareness of danger signs during pregnancy (both spontaneous and probe )	
High	2.81
Intermediate	89.13
Low	8.06
Level of awareness of danger signs during labor and delivery (both spontaneous and	

probe )	
High	0.37
Intermediate	87.16
Low	12.47
Level of awareness of danger signs during postpartum (both spontaneous and probe )	
High	10.38
Intermediate	80.75
Low	8.87
Level of awareness of danger signs in new born (both spontaneous and probe )	
High	1.59
Intermediate	89.24
Low	9.17

### **XI. Women empowerment**

Percentage of women taking own decision for health care	11.89
<b>Level of trust in health provider</b>	
High	49.79
Intermediate	32.75
Low	17.46
Percentage of women willing to contribute time in developmental activities in the village	54.29
Percentage of women willing to contribute money in developmental activities in the village	39.28

### **XII. Access to Media**

Percentage of respondents read news paper or magazine	14.56
Percentage of respondents listen to radio	28.56
Percentage of respondents watch TV	26.54
Percentage of respondents had seen drama/street play or puppet show about MNH	4.37